


Clinical Practice Guideline Blunt Liver And Spleen Injury Pediatric



- Notes:
- **No NPO**
 - **No activity restriction**
 - **No serial Hgb**
 - **No repeat CT scan**

Discharge Instructions

No activity restrictions
Expect mild increase in abdominal pain 7-10 days after injury.
Should return to baseline after 1-2 days.
Call if:

- **New, unrelenting pain**
- **Frequent light-headedness**
- **Persistent pain after 2 weeks**

	Ward Protocol	ICU Protocol
Vital signs	q2° x 4, then q4° x 24°	Continuous monitoring
Urine output	q shift	q 4°
IV access	Age appropriate size	Age appropriate size
IV fluid	Maintenance rate	Maintenance rate
Diet	Regular	Regular
Lab	Hgb on admission, and following day	Hgb on admission, 8 hrs after admission, then daily only at physician discretion
Abdominal exam	q4° x 3, and prior to discharge	q4°
Activity	Up ad lib	Up ad lib
Thresholds	Call trauma service for age adjusted tachycardia / hypotension, significant change in abdominal exam	Call trauma service for age adjusted tachycardia / hypotension, significant change in abdominal exam
Discharge criteria	To home: Hemodynamically normal x36°, no change in abdominal exam x36°	To ward: Normotensive with no tachycardia x24°, average fluid requirements